




مركز الكويت لمكافحة السرطان
Kuwait Cancer Control Center

CETUximab (every week)



Ministry of Health



* C T X - 0 0 0 0 0 - 0 1 - C N - M L *

Name:

File #:

Ht (cm):

Nationality:

Civil ID:

Wt (Kg):

Gender/Age:

DOB:

BSA (m²):

Approved indication(s): - Recurrent/Advanced stage/Metastatic Colorectal Cancer.
- Recurrent/Advanced stage/Metastatic Head and Neck Cancers.

Central line: Available NA

Allergies: NKA Yes, specify; _____

Parameters: Initiate treatment only if ANC \geq 1500; HB \geq 80; Plt \geq 75,000; CrCl $>$ 45 ml/min

Standard Protocol:

DRUG	DOSE	INSTRUCTIONS	DAYS
CETUximab	400 mg/m ² (Loading dose)	IV undiluted over 2 hr. Then, flush the IV line with 50 mL NS at end of infusion.	1st dose only
CETUximab	250 mg/m ² (maintenance dose)	IV undiluted over 60 min. Then, flush the IV line with 50 mL NS at end of infusion.	D1

To be repeated every 1 week until disease progression or intolerable toxicity.

Special instructions: Keep the patient for 1 hr observation period after completion of the 1st and 2nd cycle of CETUximab infusion.

Treatment Description:

Cycle	DATE	CETUximab	Physician	Consultant
C# __				

Post-treatment Medications: (The recommended home medications)

- Vaseline Skin care TID as long as skin is dry.
- Body lotion moisture your skin once daily as long as skin is dry.
- Fucicort cream rub the affected area with a thin film twice daily for 1 month.
- Doxycycline tab 200 mg PO daily for 2 weeks (Only if acne did not respond to topical treatment).

Comments: